



CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Roundup
READY[®] PL
HERBICIDE

with **PLANTSHIELD[®]**
 T E C H N O L O G Y

ACTIVE CONSTITUENT: 540 g/L GLYPHOSATE
 (present as the potassium salt)

GROUP M HERBICIDE

Herbicide for the control of many annual and perennial weeds in Roundup Ready Flex[®] cotton, Roundup Ready[®] canola and other situations as per the Directions for Use.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT

V 1.4

® Roundup Ready and Roundup Ready Flex are registered trademarks of Monsanto Technology LLC

APVMA Approval No.: 81975/111241

DIRECTIONS FOR USE

ROUNDUP READY FLEX[®] COTTON VARIETIES

RESTRAINTS

DO NOT disturb weeds by cultivation, sowing or grazing for six hours following treatment of annual weeds and seven days for perennial weeds.
 DO NOT use as the only method of weed control.

SITUATION – ROUNDUP READY FLEX[®] COTTON VARIETIES

IN CROP UP TO 60% BOLL OPEN STAGE

NO MORE than FOUR (4) applications¹ are permitted in crop up to 60% open stage.

Any single application in crop up to 60% open stage MUST NOT exceed 1.9 L/ha.

Total of all applications in crop must be no more than four (4) applications through all growth stages and MUST NOT exceed 7.6 L/ha.

¹ NOTE: Total of all applications of any registered glyphosate product in any one crop must not exceed 4.1 kg/ha of active constituent.

WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
Annual Ryegrass, African Turnip Weed, Annual Ground Cherry, Barnyard Grass, Bathurst Burr, Black Pigweed, Bladder Ketmia, Boggabri Weed, Button Grass, Caltrop (Yellow Vine), Camel (Afgan) Melon, Caustic Weed, Columbus Grass, Deadnettle, Liverseed Grass, Mexican Poppy, Milk (Sow) Thistle, Mintweed, Native Millet, New Zealand Spinach, Noogoora Burr, Paradoxa Grass, Pigweed (up to 25 cm diam.), Spear Thistle, Stinkgrass (Lovegrass), Sweet Summer Grass, Thornapple (Datura), Turnip Weed, Variegated Thistle, Volunteer Cereals, Volunteer Sorghum, Wild Oats, Wild/Prickly Lettuce, Wireweed	660 mL-1.9 L/ha	Rate Selection Use the lower rates on young weeds and increase to the higher rate where weeds are dense or well developed. Dense infestations of some weeds e.g. Barnyard Grass, Liverseed (Urochloa) Grass may need follow up treatments for complete control.
Climbing Buckwheat (less than 12 leaves), Couch, Johnson Grass	1.25-1.9 L/ha	Use the higher rate on plants at the flowering/seed head stage. For Johnson Grass apply to plants with a minimum of 30 cm new growth. For long term control of Couch and Johnson Grass, repeat applications will be required.
Nutgrass	1.9 L/ha followed by 1.9 L/ha	Make first application to actively growing plants when the majority of Nutgrass plants have reached at least the 6-8 leaf stage but preferably later. Allow for maximum re-emergence before retreating.

SITUATION – ROUNDUP READY FLEX® COTTON VARIETIES
IN CROP between 60% BOLL OPEN STAGE and HARVEST; QLD, NSW ONLY

NOT MORE than one (1) application.
DO NOT use on crops intended for seed production.
Application made between 60% open stage and harvest MUST NOT exceed 1.9 L/ha.
Total of all applications¹ in crop must be no more than four (4) applications through all growth stages and MUST NOT exceed 7.6 L/ha.
¹ **NOTE: Total of all applications of any registered glyphosate product in any one crop must not exceed 4.1 kg/ha of active constituent.**

WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
Bathurst Burr, Noogoora Burr, Winter annual weeds including Sowthistle/Milk Thistle	900 mL - 1.9 L/ha	Use the lower rate on light infestations of small weeds, where the crop canopy allows adequate spray coverage of the weeds. Increase to the higher rate when the crop canopy may limit spray coverage, when treating dense infestations, or when treating larger weeds. Apply alone or in tank mixtures with Dropp®. Apply when at least 60% of bolls are open and immature bolls cannot be easily cut with a sharp knife. Alternatively where the seed coat in bisected bolls is black in colour. Where a leafy canopy limits spray coverage, reduced weed control can be expected. For best results under these conditions, delay application until canopy re-opens following initial conditioning treatment.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

Tank-mixtures with other herbicides or insecticides are not recommended for over-the-top applications of this product due to the potential for reduced weed control or crop injury to result.

Tank mixes with Dropp® may be used providing the crop is 60% open and immature bolls cannot be cut with a sharp knife, alternatively where the seed coat in bisected bolls is black in colour.

WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED.

WARNING: THE APPLICATIONS RECOMMENDED ABOVE ARE FOR USE WITH IMPROVED COTTON VARIETIES THAT ARE DESIGNATED AS COTTON WITH THE ROUNDUP READY FLEX® TECHNOLOGY.

SEVERE INJURY OR DEATH OF COTTON WILL RESULT IF ANY COTTON VARIETIES NOT PROPERLY DESIGNATED AS HAVING THE ROUNDUP READY FLEX® TECHNOLOGY ARE SPRAYED WITH THIS PRODUCT.

EXTREME CARE MUST BE TAKEN TO AVOID CONTACT WITH CROPS OR DESIRABLE PLANTS WITH THE ROUNDUP READY® TECHNOLOGY, OR WITH NATIVE VEGETATION, SINCE SEVERE INJURY OR DESTRUCTION WILL RESULT.

Note: This label applies to the use of Roundup Ready® PL Herbicide on Roundup Ready Flex® cotton varieties, including Roundup Ready Flex®/Bollgard II® and Roundup Ready Flex®/Bollgard® 3 cotton varieties.

ROUNDUP READY® CANOLA VARIETIES

RESTRAINTS

DO NOT use as the only method of weed control if glyphosate resistant weeds are suspected or present.

CROP SAFETY

Applications may be made in Roundup Ready® canola varieties from crop emergence to the 6 leaf stage (prior to bud formation). Sequential applications must be at least 14 days apart and canola must have incremental growth of at least 2 new leaves between applications.

Some short-term, visual yellowing may occur when Roundup Ready® PL Herbicide is applied. This effect is temporary and will not influence crop growth or yield.

No additional surfactant is required for use in Roundup Ready® canola varieties.

Roundup Ready® PL Herbicide should be applied alone or with a 300 g/L clopyralid aqueous solution product, a 417 g/L liquid ammonium sulfate product, Asound® Duo and Nufarm Dimethoate. Other tank mixes are not recommended for over-the-top applications of this product due to the potential for reduced weed control or crop injury to result. A 417 g/L liquid ammonium sulfate product may increase the performance of this product on annual and perennial weeds, particularly under hard water conditions (high levels of calcium, magnesium or bicarbonate ions) or drought conditions.

SITUATION – ROUNDUP READY® CANOLA VARIETIES				
Before use in this situation is carried out users should consult the Roundup Ready® Canola Resistance Management Plan (RMP) which has been developed to minimise the evolution of herbicide resistance in weed populations.				
WEEDS CONTROLLED	GROWTH STAGE OF CROP	GROWTH STAGE OF WEED	RATE	CRITICAL COMMENTS
Annual Ryegrass, Barley Grass, Brome Grass, Canary Grass, Capeweed, Patersons Curse, Saffron Thistle, Scotch Thistle, Silver Grass, Spear Thistle, Variegated Thistle, Volunteer Cereals, Wild Mustard, Wild Oats, Wild Radish, Wild Turnip, Winter Grass	Crop emergence to 6 leaf (prior to bud formation).	For grass weeds and volunteer cereals: 1 leaf to mid-tillering. For volunteer plants and/or broadleaf weeds: 1 true leaf to 8 leaves.	1.15 L/ha	Up to 2 applications only may be made in any one crop. Each application must be 1.15 L/ha. Repeat applications may be required if a second flush of weeds germinates but do not apply after the 6-leaf stage of the crop. For sequential applications, applications must be at least 14 days apart and the canola crop must have incremental growth of two leaves between applications. The canola crop must have not advanced beyond the latest recommended growth stage (i.e. 6 leaf). Ensure broadleaf weeds have at least one true leaf, and grasses two leaves before application.
Weeds as above plus, Field Peas, Lupins, Sub-Clover, Annual Medic, Lentils, Chick Peas	Crop emergence to 6 leaf (prior to bud formation). Two applications required.		1.15 L/ha	Two applications of Roundup Ready® PL Herbicide provide higher levels of control than a single application.
Weeds as above plus, Faba Beans, Field Peas, Chick Peas, Lupins, Lentils, Sub. Clover, Annual Medic, Vetch	2 to 6 leaf (prior to bud formation). One or two applications.		1.15 L/ha + 150-300 mL/ha of a 300 g/L clopyralid aqueous concentrate product	Use the higher rate of a 300 g/L clopyralid aqueous concentrate product in situations of high weed population, large weed size, and/or conditions of environmental stress (dry, frost etc). Varying levels of control can be experienced between different varieties of these species. Total application of a 300 g/L clopyralid aqueous concentrate product should not exceed 300 mL/ha in the one season. Application of this tank mix at the first spray timing is recommended.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD:

Harvest: NOT REQUIRED WHEN USED AS DIRECTED.

Grazing: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

WARNING: THE APPLICATIONS RECOMMENDED ABOVE ARE FOR USE ONLY WITH IMPROVED CANOLA VARIETIES THAT ARE DESIGNATED AS CANOLA WITH THE ROUNDUP READY® TECHNOLOGY.

SEVERE INJURY OR DEATH OF CANOLA WILL RESULT IF ANY CANOLA VARIETIES NOT PROPERLY DESIGNATED AS HAVING THE ROUNDUP READY® TECHNOLOGY ARE SPRAYED WITH THIS PRODUCT.

EXTREME CARE MUST BE TAKEN TO AVOID CONTACT WITH CROPS OR DESIRABLE PLANTS WITHOUT THE ROUNDUP READY® TECHNOLOGY, OR WITH NATIVE VEGETATION, SINCE SEVERE INJURY OR DESTRUCTION WILL RESULT.

**DIRECTIONS FOR USE – GENERAL USE SITUATIONS**

ALL STATES (EXCEPT WHERE NOTED)

SITUATION	CRITICAL COMMENTS READ APPLICATION CHECKLIST BEFORE USING
GENERAL WEED CONTROL in Domestic areas (Home gardens), Commercial, Industrial and Public Service areas, Agricultural buildings and other farm situations. For specific weeds refer to the appropriate Weeds Controlled table.	For the control of many grasses and broadleaf weeds. RATE: 7 mL/L water Apply when weeds are actively growing. Apply to ensure complete and uniform wetting of foliage. Visible symptoms may take from 3 to 7 days to develop.
NON - AGRICULTURAL AREAS Around buildings, Commercial and industrial areas, Domestic and Public Service areas, Right-of ways.	Roundup Ready® PL Herbicide does not provide residual weed control. For residual control of annual weeds, Roundup Ready® PL Herbicide may be tank mixed with certain residual herbicides. See Tank Mixtures/Herbicides.
AGRICULTURAL AREAS	Roundup Ready® PL Herbicide may be used for control of annual and perennial weeds as directed, in agricultural land prior to sowing of any edible or non-edible crop, but not prior to transplanting tomato seedlings.
DRY DRAINS AND CHANNELS, DRY MARGIN OF DAMS, LAKES AND STREAM SITUATION	DO NOT apply to weeds growing in or over water. DO NOT spray across open bodies of water, and do not allow spray to enter water. DO NOT allow water to return to dry channels and drains within 4 days of application.
FORESTS	Roundup Ready® PL Herbicide may be used prior to establishment of nurseries, for site preparation prior to planting and amongst established trees using a directed or shielded spray. DO NOT allow spray or spray drift to contact foliage or green bark of desirable trees, since severe injury may result.
COTTON Shielded sprayers, QLD & NSW only. For cotton with Roundup Ready® technology see Directions for Use – Roundup Ready Flex® Cotton as appropriate.	SHIELDED SPRAYERS: Apply Roundup Ready® PL Herbicide to weeds growing between crop rows using a shielded sprayer. Refer to the Weeds Controlled tables for rates of application. DO NOT apply in crop less than 20 cm high. DO NOT allow spray or spray drift to contact any part of the cotton plant as severe injury or destruction may result.
TREE VINE CROPS Vineyards, Berries and other Small Fruits (excluding Strawberry), Citrus Fruits, Tropical and Sub-Tropical Fruits, Pome Fruits, Stone Fruits, Tree Nuts, Duboisia, Hops, Tea.	Apply as a directed or shielded spray. DO NOT apply as a spray near trees or vines less than 3 years old unless they are effectively shielded from spray and spray drift. Citrus Fruit, Nuts, Olives, Pome Fruit & Vineyards: DO NOT allow spray or spray drift to contact green bark or stems, canes, laterals, suckers, fresh wounds, foliage or fruit. Tea: Apply a maximum of 2.5 L /ha by shielded boom or directed off-centre nozzle or 4 mL /L by directed handgun or knapsack to avoid application to the crop. All other crops: DO NOT allow spray or spray drift to contact any part of the plant including the trunk. CAUTION Where split bark on Kiwifruit and green stems on Pawpaw occur, extreme care is required. For residual control of annual weeds, Roundup Ready® PL Herbicide may be tank mixed with compatible herbicides which are labeled for use in the above crops. See Tank Mixtures/Herbicides for directions.
PASTURE	DIRECTED (SPOT) APPLICATION: Roundup Ready® PL Herbicide is non-selective and may damage or kill any plant in the sprayed area. Re-treatment and/or pasture improvement may be necessary to restrict seedling re-establishment. BOOM APPLICATION: Roundup Ready® PL Herbicide may be used to suppress or kill existing pasture species prior to re-seeding or establishment of other crops. Where spot application is undertaken, grazing stock need not be removed. CAUTION Certain plants may be naturally toxic to stock. Where known toxic plants are present DO NOT allow stock to graze until complete browning of treated plants has occurred.
ONIONS Post-plant, pre-emergence application Tas only.	For control of annual weeds and suppression of perennial weeds, including Rope Twitch, apply Roundup Ready® PL Herbicide at 670 mL–2.0 L /ha post-sowing and at least 7 days before crop is due to emerge. DO NOT apply to emerging onion plants as severe injury will result. Use the lower rate on small, actively growing annual weeds. Increase to the higher rate for larger annual weeds (over 15 cm tall) and for suppression of perennial weeds.

**ANNUAL WEEDS – REGISTRATION IN ALL STATES/TERRITORIES UNLESS OTHERWISE SPECIFIED**

WEEDS CONTROLLED	BOOM RATE	HANDGUN/ KNAPSACK	CRITICAL COMMENTS
Annual Ryegrass, Amaranth, Barley Grass, Barnyard Grass, Bent Grass, Brome Grass, Caltrop, Canary Grass, Capeweed, Cereals, Chickweed, Cobbler's Peg, Deadnettle, Doublegee, Fumitory, Ground Cherry, Hedge Mustard, Hoary Cress, Lesser Swinecress, Liverseed Grass, Mintweed, Noogoora Burr, Paradoxa Grass, Paterson's Curse, Pigweed, Potato Weed, Saffron Thistle, Silver Grass, Sowthistle, Spear Thistle, Spiny Burrgrass, Spurge, Thornapple, Wild Oats, Wild Turnip, Winter Grass, Variegated Thistle	1.3–2 L /ha	4–7 mL /L	Apply to weeds whenever they are not subject to stress due to drought or frost. Use higher rate on weeds over 15 cm in height or diameter or where dense weed cover limits spray coverage. Use higher spot spraying rate when applying less than 5 L spray per 100 m ² . Roundup Ready® PL Herbicide does not provide residual weed control. Repeat treatments may be necessary to control later germinating weeds. For residual control of annual weeds Roundup Ready® PL Herbicide may be tank-mixed with certain residual herbicides. See Tank Mixtures in the General Instructions for directions. DO NOT use an atrazine tank mix for control of Barnyard Grass of Liverseed Grass.

PERENNIAL WEED – REGISTRATION IN ALL STATES/TERRITORIES UNLESS OTHERWISE SPECIFIED

WEEDS CONTROLLED	BOOM RATE	HANDGUN/ KNAPSACK	CRITICAL COMMENTS
Artichoke Thistle, African Lovegrass, Bent Grass, Carpet Grass, Cocksfoot, Flatweed, Johnson Grass, Kangaroo Grass, Kikuyu, Nutgrass (<i>Cyperus rotundus</i>), Paspalum, Phalaris, Plantain, Prairie Grass, QLD Blue Grass, Redleg Grass, Rhodes Grass, Rope Twitch, Sorrel, Soursob, #Tall Sedge, Yorkshire Fog	1.9–3.8 mL /ha	7 mL /L	Control of established perennials is best obtained when plants are at the seedhead stage. (Early flower flatweed). In general best control of winter growing perennials is obtained with application during winter-spring. Best control of summer growing perennials is obtained with application late summer and autumn. For Nutgrass in cultivated situations apply sequential treatments when Nutgrass has a minimum of 6-8 leaves. Use the higher rate in uncultivated situations. For Rhodes Grass, Rope Twitch, Prarie Grass, Qld Blue Grass, Johnson Grass, Kangaroo Grass, Kikuyu, Redleg Grass, Paspalum and Sorrel, use the higher boom rate only.
Blady Grass, Bracken, Couch, #Cumbungi, #Glyceria, Guinea Grass, #Paragrass, Silver Nightshade, #Watercouch, #See Dry Drains and Channel Use situation	5.7 mL /ha	9 mL /L	For Bracken add Pulse® at 200 mL/100 L spray mix. Best control of couch in WA and SA is obtained with spring treatment. Most effective control of Couch in eastern states is obtained with summer and autumn treatments. In cultivated situations use sequential treatments of 1.9–3.8 L/ha for control.

WOODY WEEDS – REGISTRATION IN ALL STATES/TERRITORIES UNLESS OTHERWISE SPECIFIED

WEEDS CONTROLLED	HANDGUN/ KNAPSACK	CRITICAL COMMENTS
Bamboo, Bitou Bush, Boneseed, Boxthorn, Croftonweed, Gorse, Groundsel Bush, Lantana, Mistflower	7 mL /L	Apply to actively growing plants, DO NOT apply to drought stressed plants. Further treatment may be necessary to restrict seedling reestablishment. Bamboo: apply when foliage/regrowth is 1–2 m tall. Bitou Bush/Boneseed, best results are achieved when treated at peak flowering during Winter. Groundsel Bush: DO NOT apply in Winter. Gorse: Always add Pulse® at 200 mL/100 L of spray mix, use higher rate only. Lantana: Addition of Pulse® (200 mL/100 L) may improve control. Boxthorn, Gorse, Lantana: Removal of bushes (after complete brownout), pasture improvement or further treatments are recommended to control seedlings and/or regrowth.
Blackberry, Chinese Scrub, <i>Eucalyptus</i> spp. (seedlings <2m), Hawthorn, Pampas Grass, Sifton Bush, Sweet Briar, Willow (<2m)	7-9 mL /L	Apply to actively growing plants. Removal of bushes (after complete brownout), pasture improvement or further treatments are recommended to control seedlings and/or regrowth. Blackberry: Apply from flowering to leaf fall. In Tasmania, DO NOT treat bushes bearing mature fruit. Chinese Scrub: Use higher rate on bushes greater than 1 m. <i>Eucalyptus</i> spp: Add Pulse® at 200 mL/100 L of spray mix. Hawthorn: Apply from flowering to leaf fall. Pampas Grass: Allow regrowth to reach 1m, best results – apply after flowering. Sifton Bush: Use higher rates on bushes greater than 1 m. Sweet Briar: Apply from late flowering to leaf fall, use 1.0–1.35 L/100 L, and 150–190 mL /15 L, use higher rates on bushes greater than 1.5 m

CONSERVATION TILLAGE

RESTRAINTS

To ensure herbicide absorption, DO NOT disturb weeds by cultivation, sowing or grazing for 1 day after treatment of annual weeds and 7 days for perennial weeds, except where noted.

SITUATION	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
SOUTHERN AUSTRALIA Prior to sowing a crop or pasture with FULL SOIL DISTURBANCE by cultivation or sowing with a tyned implement.	Barley Grass, Brome Grass, Volunteer Cereals, Wild Oats	340–680 mL /ha pre-tillering 680–850 mL /ha post-tillering	Treat only actively growing weeds not under stress from low moisture, frost, cold, disease or waterlogging. If heavy grazing has occurred, allow regrowth to 6-8 cm before spraying and use the higher rate. Rate Selection: Increase to higher rates late in the season or when treating under cold/overcast conditions. Full disturbance with cultivation or sowing with a tyned implement may start one day after treatment (7 days if Dock, Phalaris, Skeleton Weed, Soursob or Sorrel are present) and should occur within 21 days after treatment. Where cultivation or sowing does not occur within 21 days, new weed growth may require further treatment. When treating light infestations of seedling annual grasses (pre-tillering) and annual broadleaved weeds (less than 8cm dia/height), cultivation or sowing may start 6 hours after treatment and should occur within 21 days. Crop Establishment: Sowing should not proceed until conditions allow the formation of a satisfactory seedbed. See Crop Establishment for directions. Annual Ryegrass, Silver Grass and Perennial Grasses
	Annual Phalaris (Canary grass), Annual Ryegrass, Silver Grass, Winter Grass	680–850 mL /ha pre-tillering 850 mL–1.0 L /ha post-tillering	Addition of Wetter TX, 200 mL/100 L spray solution, may improve control. When treating dense infestation of Silver Grass, use nozzles designed to give a COARSE spray quality (ASAE S572) and a spray volume of 70 m L/ha or more is recommended to improve plant spray coverage. Good coverage of Silver Grass is critical for control. Tank Mixtures: For improved control of Clover add a 500 g/L dicamba aqueous concentrate product. Read and follow all label directions, restraints, plantback periods, withholding periods, regional use restrictions and safety directions for the tank mix products. See Tank Mixtures for directions.
	Calomba Daisy, Capeweed, Doublegee/Spiny Emex	340–680 mL /ha less than 8 cm diam/ height 680 mL–1.0 L/ha greater than 8 cm diam/ height	Perennial Weeds: For Perennial phalaris, Soursob, Skeleton Weed and Sorrel, Roundup Ready® PL Herbicide will provide knockdown, seasonal suppression and reduction in treated plant numbers.
	Amsinckia, Fumitory, Paterson's Curse, Saffron Thistle, Scotch Thistle, Spear Thistle, Variegated Thistle, Volunteer Lupins, Wild Turnip	680–850 mL /ha less than 12 cm diam/ height 850 mL–1.0 L /ha greater than 12 cm diam/ height	Tasmania: Use 1.0 L /ha on annual weeds. Increase to 2.0 L /ha where perennial weeds are being treated. To control White Clover and improve control of Sorrel and Dock, add 400mL/ha of a 500 g/L dicamba aqueous concentrate product . Observe label directions and plantback periods.
	Dock (seedling)	680mL /ha - 1 L /ha	
	Perennial Phalaris, Sorrel, Soursob, Sub. Clover	1.0 L/ha	
	Skeleton weed – fully emerged rosettes NSW only		
All the above weeds TAS only	1.0–2.0 L /ha		
SOUTHERN AUSTRALIA To commence a fallow or prior to establishing a crop or pasture with an implement that gives minimal or no soil disturbance.	Barley Grass, Wild Oats, Volunteer Cereals	680 mL–1.0 L /ha	Treat only actively growing weeds not under stress from low moisture, frost, cold, disease or waterlogging. If heavy grazing has occurred, allow regrowth to 6–8 cm before spraying and use the higher rate. Rate Selection: Use the lower rate on young weeds or where cultivation is to follow within 21 days; increase to the higher rate where grasses reach full tillering or where broadleaf weeds commence stem elongation/ budding. Increase to higher rates in spring and under cold conditions. Aerial application: Use higher rates. See Aerial Equipment .
	Brome Grass, Canary Grass, Capeweed, Variegated Thistle, Winter Grass	850 mL–1.28 L /ha	Annual Ryegrass, Silver Grass and perennial grasses: Addition of Wetter TX, 200 mL/100 L spray solution, may improve control. When treating dense infestation of Silver Grass, use nozzles designed to give a COARSE spray quality (ASAE S572) and a spray volume of 70 mL/ha or more is recommended to improve spray coverage. Good coverage of Silver Grass is critical for control. Hoary Cress: Treat from late rosette to early flowering. Soursob: Treat at tuber exhaustion.
	Annual Ryegrass, Paterson's Curse, Saffron Thistle, Scotch Thistle, Spear Thistle, Silver Grass, Wild Mustard, Wild Radish, Wild Turnip	1.0–1.28 L /ha	Couch: Use the higher rate on dense infestations. Apply sequential treatments during Summer and Autumn, with Autumn being most effective. Repeat applications will be required for full control. For improved control, use in conjunction with cultivation. Tank Mixtures: For improved control of clover add a 500 g/L dicamba aqueous concentrate product . Read and follow all label directions, restraints, plantback periods, withholding periods, regional use restrictions and safety directions for the tank mix products. See Tank Mixtures for directions. Addition of a 417 g/L liquid ammonium sulfate product, 2 L/100 L, may improve control when treating under adverse environmental conditions.
	Hoary Cress, Soursob	1 L /ha	Pasture or Crop Establishment: DO NOT sow into excessive trash. Excessive plant residues may be removed by grazing after treatment. Grazing may commence one day after treatment of annual weeds (small) and 7 days for perennial weeds. Delay grazing for three days where annual weeds are large. Sowing may proceed when excessive trash is removed, but not sooner than one day after treatment of annual weeds and 7 days for perennial weeds. See also Crop Establishment .
	Couch	1.0–2.0 L /ha	Aerial (or Surface) Seeding: Delay seeding until trash level is completely removed by grazing and/or plant decay. When establishing pasture, ensure application of fertilizer and insecticides and follow up management is undertaken as required.
	Erodium, Plantain, Perennial-Phalaris, Sorrel, Sub. Clover, Yorkshire Fog	1.27–1.67 L /ha	
	Dock, Flatweed	1.67 L /ha	
	All the above weeds TAS only	1.0 L–2.0 L /ha	Tasmania: Use 1.0 L /ha on annual weeds. Increase to 2.0 L /ha where perennial weeds are being treated. To control White Clover and improve control of Sorrel and Dock, add 400 mL/ha of a 500 g/L dicamba aqueous concentrate product. Observe label directions and plantback periods.



SITUATION	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
SOUTHERN AUSTRALIA Pasture topping For Annual Grass, Capeweed and Calomba Daisy seed-set reduction.	Barley Grass, Brome Grass, Capeweed, Silver Grass	200–300 mL /ha	Remove stock prior to treatment to allow even regrowth. Apply to Capeweed and Annual Ryegrass at FLOWERING. For other grasses, apply from HEAD to MILKY DOUGH stage. Use higher rate for dense infestations or where Annual Ryegrass is present. Apply before signs of plants "haying off". Reduction in pasture legume population may occur as a result of treatment. DO NOT apply to Clover or Medic Crops intended for seed or hay.
	Annual Ryegrass, Calomba Daisy	300 mL /ha	
Seed-head suppression of Perennial grasses.	Bent Grass	250–420 mL /ha	Timing: Treat from late October to late November. Apply before seedheads have emerged. Use the higher rate where growth is excessive and renovation is intended the following Autumn. Follow up management: Graze hard after spraying.
Poa Tussock infested pasture For reduction of ground cover allowing pasture renovation.	Most annual weeds and suppression of Poa Tussock	2.0–2.7 L /ha	Timing: Graze heavily, then remove at least 14 days before spraying to allow fresh regrowth. Apply to actively growing plants after the autumn break but before heavy frosts (March-May). Application: Increase to the higher rate may give more effective reductions. If aerial spraying, see Aerial Equipment . Follow up management: Sowing may start from 14 days after spraying. It is essential that correct follow up pasture establishment and management occurs after treatment. Spot treatment will limit re-infestation.
Serrated Tussock For control/suppression prior to establishing crops or improved pasture species NSW, Vic, Tas only.	Serrated Tussock	2.7–4 L /ha	Apply to actively growing and stress free plants. Best results May to October. Application: Boom spray volume of 70L/ha or more is recommended to improve plant coverage. Also see Aerial Equipment . Surfactants: Addition of 200mL of Wetter TX to 100L of spraying solution may improve control of Serrated Tussock. Site Preparation: Burning of Serrated Tussock 10-12 months before spraying or slashing / heavy grazing (cell grazing) 2 weeks before spraying is essential for good results (Note: Serrated Tussock is almost indigestible and prolonged exposure can lead to starvation and death of stock). Rates: Use lower rate on Serrated Tussock regrowth after burning (no residual dead foliage). Use higher rate on Serrated Tussock that has been slashed or grazed (may contain some residual dead foliage).
Serrated Tussock For prevention of seed head emergence and seed formation	Serrated Tussock	460–900 mL/ha	Apply to actively growing and stress free plants. Best results obtained during mid September – mid October. Apply prior to any seed head emergence. Also see Aerial Equipment . Surfactants: Addition of Wetter TX, 200 mL/100 L spray solution may improve results. Rates: The lower rates will be less damaging to desirable pasture species. If seed head emergence is imminent then higher rates will give better results.



SITUATION	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
NORTHERN AUSTRALIA In fallow or prior to planting a crop. QLD, NSW only	Annual Phalaris, (Canary Grass), Barley Grass, Volunteer Cereals, Wild Oats	340–680 mL /ha	<p>Treat only actively growing weeds not under stress from low moisture, frost, cold, disease or waterlogging. If heavy grazing has occurred, allow regrowth to 6-8cm before spraying and use the higher rate. Note that under summer (hot) conditions, dense infestations of Barnyard Grass and Liverseed Grass may require follow up treatment for complete control. Enhanced control of Barnyard Grass and Liverseed Grass may require follow up treatment for complete control. In winter (cold) conditions symptoms on Deadnettle may be slow to develop.</p> <p>Rate Selection: Use the lower rates on young weeds; increase to the higher rate where grasses reach full tillering or where broadleaf weeds reach stem elongation/ budding. At more advanced stages of growth certain broadleaf weeds require a higher rate range or the addition of a 680 g/L 2,4-D ethyl hexyl ester emulsifiable concentrate product.</p> <p>Crop Establishment: Sowing should not proceed until conditions allow the formation of a satisfactory seedbed. See Crop Establishment for directions.</p> <p>Tank Mixtures: Read and follow all label directions, restraints plant-back and withholding periods, regional use restrictions and safety directions for the tank mix products. DO NOT tank mix with atrazine when spraying Barnyard Grass or Liverseed Grass.</p> <p>Aerial Application: For instructions on aerial application under hot conditions see Aerial Equipment. DO NOT apply by aircraft when ambient temperature is above 30°C.</p>
	Barnyard Grass, Button Grass, Columbus Grass (seedling), Liverseed Grass, Native Millet, Stinkgrass (Lovegrass), Volunteer Sorghum	680 mL–1.27 L /ha	
	Australian Bluebell (QLD only), Cudweed, Fumitory, Mexican Poppy, New Zealand Spinach, Saffron Thistle, Spear Thistle, Spurge, Stinking Goosefoot	680 mL–1.0 L /ha	
	Black (Giant) Pigweed, Boggabri Weed, Caltrop (Yellow Vine), Indian Hedge Mustard, Mintweed, Summer Grass	340–680 mL /ha up to 5 true leaves or 3 cm dia/height 680 mL–1.0 L /ha greater than 5 true leaves or 3 cm dia/height	
	African Turnip Weed, Deadnettle, Sweet Summer Grass, Variegated Thistle, Volunteer Sunflower	510–680 mL /ha up to 5 true leaves or 3 cm dia/height 680 mL–1.27 L /ha greater than 5 true leaves or 3 cm dia/height	
	Annual Ground Cherry (Gooseberry), Bladder Ketmia, Camel Melon, False Castor Oil Plant (Thornapple), Noogoora Burr, Turnip Weed, Wild Lettuce, Wild Turnip, Wireweed	680 mL –1.0 L /ha prior to stem elongation/ budding. After stem elongation/ budding use 340 mL–1.0 L /ha plus 1.1–1.7 L/ha Surpass 475 or 1.0–1.27 L/ha of Roundup Ready® PL Herbicide alone	
	Pigweed	680 mL–1.27 L /ha	
	Sowthistle, Milk Thistle	510–680 mL /ha rosettes up to 3 cm dia 680 mL–1.27 L /ha greater than 3cm dia.	
Couch	1.0–2.0 L /ha	Use the higher rate for dense infestations. Apply sequential treatments during Summer and Autumn, with Autumn being most effective. Repeat applications will be required for full control. For improved control use in conjunction with cultivation.	
Johnson Grass	1.27–2.0 L /ha	Use the higher rate on plants approaching seedhead stage. Apply to plants with a minimum of 30 cm new growth. Sequential treatments will be required for long term control.	
Nutgrass	2.0 L + 2.0 L /ha	Make first application to actively growing plants when at least 20% have reached the head stage (normally about Feb). After allowing maximum re-emergence to occur (normally 6-8 weeks), it is essential to make a second application. Note: Follow up treatments should be made as part of a Nutgrass control program.	



SITUATION	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
SORGHUM CONTROL (pre-harvest) QLD, NSW only.	Sorghum (Grain Sorghum) - DO NOT apply to varieties intended for seed production or varieties prone to lodging	1.0–1.27 L/ha	DO NOT apply if crop is under stress from low moisture, frost, cold or water logging. Apply when grain moisture is less than 25%. Use the higher rate where the crop has produced significant number of late tillers or where following crops will be established without further treatment. Pre-harvest treatments may increase the likelihood of crop lodging. Apply post-harvest treatments to previously slashed/grazed stubble when least 20 cm of new growth has occurred. Use the higher rate on standing stubble or where re-growth from slashed Sorghum has advanced beyond 50cm in height. CAUTION Sorghum may be naturally toxic to stock.
SORGHUM CONTROL (post-harvest) QLD, NSW only.	Sorghum stubble (grain-sorghum)	680 mL–1.0 L/ha for fresh regrowth from slashed stubble. 1.0 L–1.27 L/ha for standing stubble if sufficiently green and for fresh spring regrowth	
SUGARCANE Ratoon Spray out QLD, NSW only.	Sugarcane ratoon regrowth	2.7–6 L /ha	APPLY UNDER GOOD GROWING CONDITIONS ONLY to actively growing ratoons 60–120 cm tall. DO NOT apply if plants are under stress from low moisture or waterlogging. Use the lower rate for suppression or where cultivation is to follow. Use the higher rate for control.
RICE Direct drilling NSW only.	Annual Phalaris (Canary Grass), Annual Ryegrass, Barley Grass, Burr Medic, Sub. Clover, Winter Grass	680–840 mL /ha	Roundup Ready® PL Herbicide is less effective in drought-stressed plants. In drought conditions a pre-watering prior to spraying is recommended. In grazed situations, if heavy grazing has occurred allow regrowth to 6–8 cm before spraying. Annual Ryegrass: Add Wetter TX at 200 mL/100 L of spray solution and where dominant, use the higher rate. Sowing: Direct drilling may take place 1–14 days after spraying. Roundup Ready® PL Herbicide does not provide residual weed control. Permanent water and approved selective herbicides should be used to provide continuing control of weeds.
PRE-HARVEST APPLICATION to reduce viable seed set of weeds in: Field Peas (<i>Pisum sativum</i>) Faba Beans (<i>Vicia faba</i>)	Annual Ryegrass (<i>Lolium rigidum</i>)	320–680 mL /ha	Use lower rate if Ryegrass is flowering and higher rate if Ryegrass is at milky dough stage. Application should be made at or after crop maturity. Application before this time may significantly reduce yields (in practice losses in excess of 25% can occur). Apply when the average seed moisture content is below 30%. For Faba Beans, this is indicated by the pods going black, and for Field Peas by the pods going yellow. DO NOT harvest within 7 days after application. DO NOT use on crops intended for seed or sprouting. Glyphosate resistant biotypes have been detected in Australia. If glyphosate resistant weeds are known to be present, apply an additional method of control.
PRE-HARVEST APPLICATION as harvest aid and weed control: Wheat (<i>Triticum aestivum</i>)	Annual weeds	900 mL–1.8 L /ha	Apply to mature crop from late dough stage (28% moisture) onwards. The higher rate will be required when crops are heavy and leaf shading effects may occur. DO NOT harvest within 7 days after application. DO NOT use on crops intended for seed or sprouting. Where Wheat is grown in rotation with any herbicide tolerant crop, management should be consistent with implementation of any management plan for herbicide tolerant crops.
PRE-HARVEST APPLICATION To desiccate a crop as a harvest aid and weed control Adzuki Beans, Chickpeas, Cowpea, Faba Beans, Field Peas, Lentils, Mungbeans, Soybean (Application to crops intended for seed production or for sprouting may reduce germination percentage to commercially unacceptable levels.)	Annual Weeds	680 mL–1.8 L /ha	Apply with boom or by air. Use higher rates where crops or weeds are dense and where faster desiccation is required. Application should be made at or after crop maturity: Chickpeas and Lentils: Apply when physiologically mature and less than 15% green pods. Soybean: Apply only after seed pods have lost all green colour and 80-90% of leaves have dropped. Use only on soybean crops grown for crushing. Mungbeans/ Adzuki and Cowpea: Apply to mature crops when pods are brown/black. Field peas: Apply when seeds turn yellow and average seed moisture content is below 30%. Faba Beans: Apply when pods turn black and average seed moisture content is below 30%. DO NOT harvest within 7 days of application. Speed of crop desiccation is dependent on crop stage, growing conditions and weather conditions during and after application. This use should be part of an Integrated Weed Management strategy which incorporates herbicides with different modes of action and alternative cultural weed control practices.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIOD

WHEAT AND LEGUMES: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.

ALL OTHER USES: NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

PRODUCT INFORMATION

Roundup Ready® PL Herbicide is a non-volatile, non selective water soluble liquid herbicide for the control of annual and perennial grasses and broadleaf weeds in Roundup Ready Flex® cotton, Roundup Ready® canola and certain other situations. Roundup Ready® PL Herbicide is absorbed by plant foliage and green stems. Roundup Ready® PL Herbicide is inactivated on clay and organic matter in soil and does not provide residual weed control. Roundup Ready® PL Herbicide moves throughout the weed from the point of contact to and into the root system. Initial visible effects on annual weeds take 3-7 days but may not be noticeable for 2 to 3 weeks under cool cloudy conditions or on some perennial weeds. Roundup Ready® PL Herbicide will not control Roundup Ready Flex® cotton or Roundup Ready® canola volunteers at any leaf stage.

RESISTANT WEEDS WARNING

Roundup Ready® PL Herbicide is a member of the Glycines group of herbicides. Roundup Ready® PL Herbicide has the inhibition of EPSP synthase mode of action. For weed resistance management Roundup Ready® PL Herbicide is a Group M herbicide. Some naturally occurring weed biotypes resistant to Roundup Ready® PL Herbicide and other Group M herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Roundup Ready® PL Herbicide or other Group M herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Monsanto accepts no liability for any losses that may result from the failure of Roundup Ready® PL Herbicide to control resistant weeds.

Users of Roundup Ready® PL Herbicide over Roundup Ready Flex® cotton or Roundup Ready® canola must implement practices that minimise the development of resistance in treated weeds. Minimising this risk may best be achieved by following the integrated weed management strategy guidelines summarised below:

1. Aim to enter the Roundup Ready® cropping phase of the rotation with a low weed burden.
2. Integrate as many different weed control options (chemical and cultural) as possible through all phases of the crop rotation.
3. Make every herbicide application count – use registered rates at the correct application growth stage and assess effectiveness.
4. Rotate herbicides with different modes of action throughout the crop rotation.
5. Regularly monitor the effectiveness of resistance management practices.
6. Test weed populations for herbicide resistance status as part of ongoing integrated weed management.
7. Growers should not plant Roundup Ready® crops in paddocks with populations of confirmed glyphosate resistant weeds.

It is advised that consultation on Integrated Weed Management be undertaken with an accredited agronomist or program prior to use of Roundup Ready® PL Herbicide over Roundup Ready Flex® cotton or Roundup Ready® canola.

More information on Integrated Weed Management can be found at: <http://www.glyphosateresistance.org.au> and www.weedsmart.org.au.

GROUP M HERBICIDE



As with conventional varieties, volunteer and ratoon Roundup Ready Flex® plants may occur in fallows, and non-cropping areas of a farm such as irrigation ditches, module pads, water storages, etc. These plants will not be controlled by Roundup Ready® PL Herbicide or other glyphosate (Group M) herbicides and should be controlled in both cropping and non-cropping areas. These plants are best managed with cultivation and/or the appropriate registered herbicides (see the Integrated Weed Management Strategy Guidelines above). Growers should ensure that they have an effective weed management strategy developed for the control of these weeds.

Users of Roundup Ready® PL Herbicide over Roundup Ready Flex® cotton must allow Monsanto or its agents to undertake audits or surveys as necessary to assess management by users of the development of glyphosate resistance in target weeds. Monsanto or its agents will conduct an audit or survey annually on a percentage of fields where Roundup Ready® PL Herbicide has been used over Roundup Ready Flex® cotton.

RESISTANT WEEDS REPORTING

Users of Roundup Ready® PL Herbicide and Monsanto Technology Service Providers (TSPs) are required to report any adverse events, such as suspected weed resistance, to Monsanto as soon as it is identified. Monsanto will investigate the incident and produce a report of any incidents of confirmed resistance of weeds to Roundup Ready® PL Herbicide in target weed species which are normally susceptible to this herbicide and forward the report as soon as practicable to the Australian Pesticides and Veterinary Medicines Authority. Weeds identified to have survived Roundup Ready® PL Herbicide must be controlled by and alternative strategy in order to prevent seed from setting seed.

MIXING

Roundup Ready® PL Herbicide mixes readily with water. Reduced results may occur if water is used containing suspended clay or organic matter e.g. from dams, streams and irrigation channels or high levels of calcium, magnesium or bicarbonate ions.

DO NOT mix, store or apply this product in galvanised steel or unlined steel containers or spray tanks, since a highly flammable gas mixture may be formed. Use stainless steel, aluminium, brass, copper, fibreglass, plastic or plastic lined containers or spray tanks. Spray tanks, pumps, lines and nozzles should be thoroughly cleaned with clean water following application. Ensure that the spray tank is free of any residue of other spray solutions prior to mixing. Use spray solutions promptly as a gradual loss of activity may occur over a period of days following spray preparation.

General Use and Tank Mixing Procedure

DO NOT use tank mixtures for “over the top” applications in Roundup Ready Flex® cotton unless specified in the Directions for Use. Consider carefully any plant back periods to cotton or other crops.

1. Fill the spray tank 1/3 or 1/2 full with clean water and start agitation.
2. Where a crystalline ammonium sulfate is recommended, add 1 kg/100 L spray solution into the tank and mix thoroughly.
3. Add recommended herbicide/additive to the spray tank and mix thoroughly.
4. Add Roundup Ready® PL Herbicide. Mix thoroughly and continue water addition.
5. Always maintain adequate agitation during application and use the tank-mix promptly.

Clean all equipment after use by washing thoroughly with water or recommended decontaminant.

**TANK MIXTURES**

IMPORTANT: NOT FOR USE OVER THE TOP OF ROUNDUP READY FLEX® COTTON UNLESS SPECIFIED IN THE DIRECTIONS FOR USE.

HERBICIDES

Ally® Herbicide, Affinity® Force Herbicide, 300 g/L clopyralid aqueous concentrate product, 600 g/kg metsulfuron-methyl water dispersible granule, 500 g/L tri-allate, 400 g/L fluroxypyr, Eclipse®, 100 SC Herbicide, 680 g/L 2,4-D ethyl hexyl ester emulsifiable concentrate, Express® Herbicide, Flame® Herbicide, Garlon® 600 Herbicide, Hammer® 400 EC Herbicide, 600 g/L triclopyr, 500 g/L dicamba aqueous concentrate product, Lontrel® 750 SG Herbicide, 570 g/L MCPA LVE, Monza®, 750 g/kg triasulfuron, 600 g/L atrazine soluble concentrate, 900 g/kg atrazine water dispersible granule, 500 g/L simazine suspension concentration, 900 g/kg simazine water dispersible granule, Stomp® 440 Herbicide, 240 g/L oxyfluorfen, Surflan 500 Flowable Herbicide, 475 g/L 2,4-D amine, 480 g/L trifluralin and Yield® 250 EC Herbicide. Two-way mixes are chemically and biologically compatible, other mixes (three-way) have not been tested.

The addition of Striker® at 75 mL/ha to recommended rates of Roundup Ready® PL Herbicide prior to planting winter cereals or cotton will improve knockdown of certain weeds.

INSECTICIDES

This product is compatible with the following insecticides. 100 g/L alpha-cypermethrin, 400 g/L dimethoate, Imidan®, Karate® Zeon Insecticide, O-Mat® 290 SL Insecticide, Lorsban® 500 EC Insecticide, Sumithion ULV and emulsifiable concentrates of fenitrothion. Two-way mixes are chemically and biologically compatible, other mixes (three-way) have not been tested. Other insecticides have not been tested.

SURFACTANT ADDITION

Additional surfactant is not required except where the rate of Roundup Ready® PL Herbicide is less than 7 mL/L (eg. 700 mL/100 L water) when applied by boom.

ADDITIVES

A 417 g/L liquid ammonium sulfate product may be used as an adjuvant to alleviate the adverse effects of high levels of calcium, magnesium and bicarbonate ions in water.

RATE: 2 L per 100 litres spray solution.

Pulse® Penetrant

RATE: 20 mL/100 L spray solution.

Add when treating Bracken and many woody weeds.

Wetter TX Surfactant

RATE: 20 mL/100 L spray solution.

Add when treating Annual ryegrass in spring (from beginning August to end October). Wetter TX is not a general purpose surfactant and should be used only where recommended.

APPLICATION**Ground Boom**

For broadcast (over-the-top) application, a spray volume of 50–80 litres per sprayed hectare is recommended for optimum performance. Nozzles and pressure settings must be selected to deliver a minimum of COARSE spray quality (American Society of Agricultural Engineers (ASAE) S572) at the target. Depending on prevailing temperature, relative humidity, delta T, wind speed, travel speed and boom height the spray quality produced at the nozzles may need to be coarser than this. In sensitive areas avoid using nozzles and/or pressure settings that produce a VERY FINE to MEDIUM spray quality, as these droplets are more prone to drift off-target.

Directed / Shielded Ground Application Equipment

Equipment should be used which directs the spray plume to the base of the cotton plants minimising contact with the foliage. Total application volume of 80 L/ha should be used. Select nozzle types that produce a minimum COARSE spray quality (ASAE S572). Be aware of operational factors such as ground speed, nozzle height and row integrity. Monitor the application using water sensitive paper if uncertain. Monitor environmental conditions that may influence off target droplet movement such as temperature, relative humidity and wind speed.

High Volume Application

(e.g. Knapsack/Handgun Equipment.) The dilution rate is given as g/litre e.g. 7 mL Roundup Ready® PL Herbicide per 1 L of water. This is equal to 100 mL Roundup Ready® PL Herbicide per 15 L of water or 640 mL per 100 L of water. Adjust equipment to achieve an even spray pattern with a minimum of a COARSE spray quality at the target. Apply to ensure complete and uniform wetting of all foliage.

Aerial Equipment

When applying Roundup Ready® PL Herbicide by air over the top (OTT) of Roundup Ready Flex® cotton nozzles and pressure settings must be selected to deliver a minimum of a COARSE spray quality (ASAE S572) at the target. Depending on prevailing temperature, relative humidity, delta T, wind speed, travel speed and boom height the spray quality produced at the nozzles may need to be coarser than this. In sensitive areas avoid using nozzles and/or pressure settings that produced a VERY FINE to MEDIUM spray quality as these droplets are more prone to drift off-target. A minimum total application volume of 40 L per hectare needs to be used.

DO NOT apply Roundup Ready® PL Herbicide by aircraft at temperatures above 30°C. Avoid application when relative humidity falls below 35%.

DO NOT apply during low-level inversion conditions, when winds are gusty or under any other conditions which favour drift. Drift may cause damage to any vegetation contacted to which treatment is not intended. To prevent injury to adjacent desirable vegetation, appropriate buffer zones must be maintained.

PREVAILING ENVIRONMENTAL CONDITIONS MUST BE CONSIDERED.

ANY AERIAL APPLICATION TO COTTON SHOULD BE DONE IN ACCORDANCE WITH THE AUSTRALIAN COTTON INDUSTRY'S BEST MANAGEMENT PRACTICES MANUAL.

AVOID DRIFT - EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS WHICH DO NOT CONTAIN THE APPROPRIATE ROUNDUP READY® TECHNOLOGY, AND TO NATIVE VEGETATION, AND TO PREVENT CONTAMINATION OF OPEN BODIES OF WATER AND WATERWAYS.

APPLICATION CHECK LIST

- Do not treat weeds under poor growing conditions due to moisture stress, waterlogging, severe frosting, insect damage etc. Reduced performance may also occur where weeds are covered with dust or silt.
- Rain within 2 hours of application which causes run-off will require re-treatment. Rainfastness is reduced if weeds are not actively growing, under stress or conditions of low light intensity/darkness.
- Delay treatment of plants wet with dew or rain, if water droplets run off when plants are disturbed.
- Apply treatments to weeds which have at least one true leaf (broadleaf weeds) or two leaves (grasses) to provide an adequate surface area for herbicide uptake.
- Be aware of any crops that may be in the vicinity of the application that are sensitive to Roundup Ready® PL Herbicide.
- When applying Roundup Ready® PL Herbicide by air over the top of Roundup Ready Flex® cotton up to the 22nd node, nozzles and pressure settings must be selected to deliver a minimum COARSE spray quality (ASAE S572) at the target. A minimum total volume of 40 L per hectare must be used.
- If glyphosate resistant weeds are known to be present, use an alternative method of control before these weeds set seed.
- Be aware of native and other non-target vegetation in the vicinity of application, as such vegetation may be severely affected or destroyed by Roundup Ready® PL Herbicide.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS AND LIVESTOCK

Avoid contact with foliage, green stems or fruit of crops, desirable plants and trees, since severe injury or destruction may result. DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands, pasture, native vegetation or any other non-target vegetation.

PROTECTION OF WILDLIFE, FISH CRUSTACEA AND ENVIRONMENT

DO NOT contaminate dams, rivers or streams with the product or used container. DO NOT apply to weeds growing in or over water. DO NOT spray across open bodies of water.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure-rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Will irritate the eyes. May irritate the nose and throat. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When opening the container and preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist or equivalent clothing and elbow-length PVC gloves and face shield or goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. **Phone 13 11 26.**

SAFETY DATA SHEET

For further information refer to the Safety Data Sheet (SDS), which can be obtained from your supplier or the Sinochem International Australia website – www.sinochem.com.au

IMPORTANT MANUFACTURER'S NOTICE

MONSANTO DOES NOT WARRANT THE SAFETY OR PERFORMANCE OF THE ROUNDUP READY® TECHNOLOGY OR ROUNDUP READY® PL HERBICIDE WHEN SAVED COTTON OF ROUNDUP READY FLEX® COTTON ARE USED.

CAUTION: PLEASE READ THIS NOTICE BEFORE OPENING THE CONTAINER

The results obtained from using this product may be affected by factors beyond Monsanto's control, including mixing, use, climatic conditions, time of application, crop or crop stage and the possible development of resistance to the active ingredient.

Limitation of liability

With the express exception of liabilities created by the *Competition and Consumer Act 2010* (Cth)(including the Australian Consumer Law) or relevant State legislation which cannot be excluded, restricted or modified, none of Monsanto Australia Limited or any of its affiliates ("Monsanto"), or any manufacturer of any component of the product or Sinochem International Australia Pty Ltd or any of its affiliates ("Sinochem") shall be liable for any loss or damage (including consequential loss or damage), injury or death connected with, or arising out of, the product, regardless of the way in which it arises (including by way of negligence).

Remedy for failure to comply with consumer guarantees

If there has been a failure to comply with a consumer guarantee (other than a guarantee under sections 51, 52 or 53 of Australian Consumer Law or corresponding provisions in the relevant State legislation) in relation to a good which is not a good of a kind ordinarily acquired for personal, domestic or household use or consumption, the liability of Monsanto or Sinochem is limited to a replacement of the good or the supply of an equivalent good.

Distributed by:

Sinochem International Australia Limited
8/606 St Kilda Road
Melbourne Victoria 3004
Tel: +61 3 9520 8888
www.sinochem.com.au

® Roundup Ready, Roundup Ready Flex, Bollgard and Monza are registered trademarks of Monsanto Technology LLC.